

兔抗 GLYCTK 多克隆抗体

中文名称：兔抗 GLYCTK 多克隆抗体

英文名称：Anti-GLYCTK rabbit polyclonal antibody

别 名：HBEBP2; HBEBP4; HBeAgBP4A

相关类别：一抗

储 存：冷冻 (-20°C) 避光

抗 原：GLYCTK

宿 主：Rabbit

反应种属：Human Mouse Rat

标记物：Unconjugate

克隆类型：Unconjugate

技术规格

Background:	This locus encodes a member of the glycerate kinase type-2 family. The encoded enzyme catalyzes the phosphorylation of (R)-glycerate and may be involved in serine degradation and fructose metabolism. Decreased activity of the encoded enzyme may be associated with the disease D-glyceric aciduria. Alternatively spliced transcript variants have been described.
Applications:	WB, IHC, IF
Name of antibody:	GLYCTK
Immunogen:	Synthesized peptide derived from internal of human GLCTK.
Full name:	glycerate kinase

Synonyms :	HBEBP2; HBEBP4; HBeAgBP4A
SwissProt:	Q8IVS8
IHC positive control:	Human brain tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	55 kDa
WB Positive control:	NIH/3T3 cells lysate
WB Recommended dilution:	500-3000
IF positive control:	A549 cells
IF Recommend dilution:	100-500



